

Beckman-Quarles type theorems for mappings from $\mathbb{R}^n$ to $\mathbb{C}^n$

APOLONIUSZ TYSZKA

Summary. Let $\varphi_n : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$, $\varphi_n((x_1, \dots, x_n), (y_1, \dots, y_n)) = (x_1 - y_1)^2 + \dots + (x_n - y_n)^2$. We say that $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ preserves distance $d \geq 0$ if for each $x, y \in \mathbb{R}^n$ $\varphi_n(x, y) = d^2$ implies $\varphi_n(f(x), f(y)) = d^2$. Let A_n denote the set of all positive numbers d such that any map $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ that preserves unit distance preserves also distance d . Let D_n denote the set of all positive numbers d with the property: if $x, y \in \mathbb{R}^n$ and $|x - y| = d$ then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that any map $f : S_{xy} \rightarrow \mathbb{C}^n$ that preserves unit distance preserves also the distance between x and y . Obviously, $D_n \subseteq A_n$. We prove: **(1)** $A_n \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$, **(2)** for $n \geq 2$ D_n is a dense subset of $(0, \infty)$. Item **(2)** implies that each mapping f from $\mathbb{R}^n$ to $\mathbb{C}^n$ ($n \geq 2$) preserving unit distance preserves all distances, if f is continuous with respect to the product topologies on $\mathbb{R}^n$ and $\mathbb{C}^n$.

The classical Beckman-Quarles theorem states that each unit-distance preserving mapping from $\mathbb{R}^n$ to $\mathbb{R}^n$ ($n \geq 2$) is an isometry, see [1], [2], [6] and [10]. By a *complex isometry* of $\mathbb{C}^n$ we understand any map $f : \mathbb{C}^n \rightarrow \mathbb{C}^n$ of the form

$$f(z_1, z_2, \dots, z_n) = (z'_1, z'_2, \dots, z'_n)$$

where

$$z'_j = a_{0j} + a_{1j}z_1 + a_{2j}z_2 + \dots + a_{nj}z_n \quad (j = 1, 2, \dots, n),$$

the coefficients a_{ij} are complex and the matrix $\|a_{ij}\|$ ($i, j = 1, 2, \dots, n$) is orthogonal

Mathematics Subject Classification (2000). 51M05.

Keywords. Beckman-Quarles theorem, Cayley-Menger determinant.

Abbreviated title: Beckman-Quarles type theorems.

i.e. satisfies the condition

$$\sum_{j=1}^n a_{\mu j} a_{\nu j} = \delta_{\nu}^{\mu} \quad (\mu, \nu = 1, 2, \dots, n)$$

with Kronecker's delta. Let $\varphi_n : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$, $\varphi_n((x_1, \dots, x_n), (y_1, \dots, y_n)) = (x_1 - y_1)^2 + \dots + (x_n - y_n)^2$. The function φ_n is not the square of a distance because it admits values that are not real. Nevertheless, according to [4] $\varphi_n(x, y)$ is invariant under complex isometries i.e. for every complex isometry $f : \mathbb{C}^n \rightarrow \mathbb{C}^n$

$$\forall x, y \in \mathbb{C}^n \quad \varphi_n(f(x), f(y)) = \varphi_n(x, y).$$

Therefore, we say that $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ preserves distance $d \geq 0$ if for each $x, y \in \mathbb{R}^n$ $\varphi_n(x, y) = d^2$ implies $\varphi_n(f(x), f(y)) = d^2$.

By a field endomorphism of $\mathbb{C}$ we understand any map $g : \mathbb{C} \rightarrow \mathbb{C}$ satisfying:

$$\begin{aligned} \forall x, y \in \mathbb{C} \quad g(x + y) &= g(x) + g(y), \\ \forall x, y \in \mathbb{C} \quad g(x \cdot y) &= g(x) \cdot g(y), \\ g(0) &= 0, \\ g(1) &= 1. \end{aligned}$$

Bijjective endomorphisms are called automorphisms, for more information on field endomorphisms and automorphisms of $\mathbb{C}$ the reader is referred to [9] and [13]. If r is a rational number, then $g(r) = r$ for any field endomorphism $g : \mathbb{C} \rightarrow \mathbb{C}$. Proposition 1 shows that only rational numbers r have this property:

Proposition 1. If $r \in \mathbb{C}$ and r is not a rational number, then there exists a field automorphism $g : \mathbb{C} \rightarrow \mathbb{C}$ such that $g(r) \neq r$.

Proof. Note first that if E is any subfield of $\mathbb{C}$ and if g is an automorphism of E , then g can be extended to an automorphism of $\mathbb{C}$. This follows from [5, corollaire 1 on p. 109] by taking $\Omega = \mathbb{C}$ and $K = \mathbb{Q}$. Now let $r \in \mathbb{C} \setminus \mathbb{Q}$. If r is algebraic over $\mathbb{Q}$, let E be the splitting field in $\mathbb{C}$ of the minimal polynomial μ of r over $\mathbb{Q}$ and let $r' \in E$ be any other root of μ . Then there exists an automorphism g of E that sends r to r' , see for example [7, corollary 2 on p. 66]. If r is transcendental over $\mathbb{Q}$, let $E = \mathbb{Q}(r)$ and let $r' \in E$ be any other generator of E (e.g. $r' = 1/r$). Then there exists an automorphism g of E that sends r to r' . In each case, g can be extended to an automorphism of $\mathbb{C}$.

If $g : \mathbb{C} \rightarrow \mathbb{C}$ is a field endomorphism then $(g|_{\mathbb{R}}, \dots, g|_{\mathbb{R}}) : \mathbb{R}^n \rightarrow \mathbb{C}^n$ preserves all distances $\sqrt{r}$ with rational $r \geq 0$. Indeed, if $(x_1 - y_1)^2 + \dots + (x_n - y_n)^2 = (\sqrt{r})^2$ then

$$\begin{aligned}
\varphi_n((g|_{\mathbb{R}}, \dots, g|_{\mathbb{R}})(x_1, \dots, x_n), (g|_{\mathbb{R}}, \dots, g|_{\mathbb{R}})(y_1, \dots, y_n)) &= \\
\varphi_n((g(x_1), \dots, g(x_n)), (g(y_1), \dots, g(y_n))) &= \\
(g(x_1) - g(y_1))^2 + \dots + (g(x_n) - g(y_n))^2 &= \\
(g(x_1 - y_1))^2 + \dots + (g(x_n - y_n))^2 &= \\
g((x_1 - y_1)^2) + \dots + g((x_n - y_n)^2) &= \\
g((x_1 - y_1)^2 + \dots + (x_n - y_n)^2) &= g((\sqrt{r})^2) = g(r) = r = (\sqrt{r})^2.
\end{aligned}$$

Theorem 1 shows that we cannot expect a result like that of Beckman and Quarles:

Theorem 1. If $x, y \in \mathbb{R}^n$ and $|x - y|^2$ is not a rational number, then there exists $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ that does not preserve the distance between x and y although f preserves all distances $\sqrt{r}$ with rational $r \geq 0$.

Proof. There exists an isometry $I : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $I(x) = (0, 0, \dots, 0)$ and $I(y) = (|x - y|, 0, \dots, 0)$. By Proposition 1 there exists a field automorphism $g : \mathbb{C} \rightarrow \mathbb{C}$ such that $g(|x - y|^2) \neq |x - y|^2$. Thus $g(|x - y|) \neq |x - y|$ and $g(|x - y|) \neq -|x - y|$. Therefore $(g|_{\mathbb{R}}, \dots, g|_{\mathbb{R}}) : \mathbb{R}^n \rightarrow \mathbb{C}^n$ does not preserve the distance between $(0, 0, \dots, 0) \in \mathbb{R}^n$ and $(|x - y|, 0, \dots, 0) \in \mathbb{R}^n$ although $(g|_{\mathbb{R}}, \dots, g|_{\mathbb{R}})$ preserves all distances $\sqrt{r}$ with rational $r \geq 0$. Hence $f := (g|_{\mathbb{R}}, \dots, g|_{\mathbb{R}}) \circ I : \mathbb{R}^n \rightarrow \mathbb{C}^n$ does not preserve the distance between x and y although f preserves all distances $\sqrt{r}$ with rational $r \geq 0$.

Let A_n denote the set of all positive numbers d such that any map $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ that preserves unit distance preserves also distance d . By Theorem 1 $A_n \subseteq \{d > 0 : d^2 \in \mathbb{Q}\}$. Let D_n denote the set of all positive numbers d with the following property:

if $x, y \in \mathbb{R}^n$ and $|x - y| = d$ then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that any map $f : S_{xy} \rightarrow \mathbb{C}^n$ that preserves unit distance preserves also the distance between x and y .

In case of $|x - y| = 1$ we define $S_{xy} := \{x, y\}$. Thus, $1 \in D_n$. Obviously, $D_n \subseteq A_n$.

We shall study the set D_n and a unit-distance preserving mapping f from $\mathbb{R}^n$ to $\mathbb{C}^n$. We need the following technical Propositions 2-7.

Proposition 2 (cf. [3], [4]). The points $c_1 = (z_{1,1}, \dots, z_{1,n}), \dots, c_{n+1} = (z_{n+1,1}, \dots, z_{n+1,n}) \in \mathbb{C}^n$ are affinely dependent if and only if their Cayley-Menger determinant $\Delta(c_1, \dots, c_{n+1}) :=$

$$\det \begin{bmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & 0 & \varphi_n(c_1, c_2) & \dots & \varphi_n(c_1, c_{n+1}) \\ 1 & \varphi_n(c_2, c_1) & 0 & \dots & \varphi_n(c_2, c_{n+1}) \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \varphi_n(c_{n+1}, c_1) & \varphi_n(c_{n+1}, c_2) & \dots & 0 \end{bmatrix}$$

equals 0.

Proof. It follows from the equality

$$\left(\det \begin{bmatrix} z_{1,1} & z_{1,2} & \dots & z_{1,n} & 1 \\ z_{2,1} & z_{2,2} & \dots & z_{2,n} & 1 \\ \dots & \dots & \dots & \dots & \dots \\ z_{n+1,1} & z_{n+1,2} & \dots & z_{n+1,n} & 1 \end{bmatrix} \right)^2 = \frac{(-1)^{n+1}}{2^n} \cdot \Delta(c_1, \dots, c_{n+1}).$$

Proposition 3 (cf. [3], [4]). For any points $c_1, \dots, c_{n+k} \in \mathbb{C}^n$ ($k = 2, 3, 4, \dots$) their Cayley-Menger determinant equals 0 i.e. $\Delta(c_1, \dots, c_{n+k}) = 0$.

Proof. Assume that $c_1 = (z_{1,1}, \dots, z_{1,n}), \dots, c_{n+k} = (z_{n+k,1}, \dots, z_{n+k,n})$. The points $\tilde{c}_1 = (z_{1,1}, \dots, z_{1,n}, 0, \dots, 0)$, $\tilde{c}_2 = (z_{2,1}, \dots, z_{2,n}, 0, \dots, 0), \dots, \tilde{c}_{n+k} = (z_{n+k,1}, \dots, z_{n+k,n}, 0, \dots, 0) \in \mathbb{C}^{n+k-1}$ are affinely dependent. Since $\varphi_n(c_i, c_j) = \varphi_{n+k-1}(\tilde{c}_i, \tilde{c}_j)$ ($1 \leq i \leq j \leq n+k$) the Cayley-Menger determinant of points $c_1, \dots, c_{n+k}$ is equal to the Cayley-Menger determinant of points $\tilde{c}_1, \dots, \tilde{c}_{n+k}$ which equals 0 according to Proposition 2.

From Proposition 2 we obtain the following Propositions 4a and 4b.

Proposition 4a. If $d, e > 0$ with $2ne^2 \neq (n-1)d^2$ and if points $c_0, \dots, c_n \in \mathbb{C}^n$ satisfy $\varphi_n(c_i, c_j) = d^2$ for $1 \leq i < j \leq n$ and $\varphi_n(c_0, c_i) = e^2$ for $1 \leq i \leq n$, then the points $c_0, \dots, c_n$ are affinely independent.

Proposition 4b. If $d > 0$, $c_1, c_2, c_3 \in \mathbb{C}^2$ and $\varphi_2(c_1, c_2) = 3d^2$, $\varphi_2(c_1, c_3) = 11d^2$, $\varphi_2(c_2, c_3) = 25d^2$, then c_1, c_2, c_3 are affinely independent.

Proposition 5 (cf. [4] p. 127 in the real case). If points $c_0, c_1, \dots, c_n \in \mathbb{C}^n$ are affinely independent, $x, y \in \mathbb{C}^n$ and $\varphi_n(x, c_0) = \varphi_n(y, c_0)$, $\varphi_n(x, c_1) = \varphi_n(y, c_1), \dots, \varphi_n(x, c_n) = \varphi_n(y, c_n)$, then $x = y$.

Proof. Computing we obtain that the vector $\vec{xy} := [s_1, \dots, s_n]$ is perpendicular to each of the n linearly independent vectors $\vec{c_0c_i}$ ($i = 1, \dots, n$). Thus the vector $\vec{xy}$ is perpendicular to every linear combination of vectors $\vec{c_0c_i}$ ($i = 1, \dots, n$). In particular, the vector $\vec{xy} = [s_1, \dots, s_n]$ is perpendicular to the vector $[\bar{s}_1, \dots, \bar{s}_n]$, where $\bar{s}_1, \dots, \bar{s}_n$ denote numbers conjugate to the numbers $s_1, \dots, s_n$, respectively. Therefore $\vec{xy} = 0$ and the proof is completed.

Proposition 6a. The set $\left\{ \left(\sqrt{11}/5 \right)^k \cdot \left(\sqrt{3} \right)^l : k, l \in \{0, 1, 2, \dots\} \right\}$ is a dense subset of $(0, \infty)$.

Proof. It suffices to prove that $\left\{ \ln_3 \left(\left(\frac{11}{25} \right)^k \cdot 3^l \right) : k, l \in \{0, 1, 2, \dots\} \right\}$ is a dense subset of $\mathbb{R}$. Computing we obtain the set $\left\{ l + k \cdot \ln_3 \left(\frac{11}{25} \right) : k, l \in \{0, 1, 2, \dots\} \right\}$ which is a dense subset of $\mathbb{R}$ due to Kronecker's theorem ([8]), because $\ln_3 \left(\frac{11}{25} \right)$ is irrational and negative. (If there were $a, b \in \{1, 2, 3, \dots\}$ with $\ln_3 \left(\frac{11}{25} \right) = -\frac{a}{b}$, then we would have $3^a \cdot 11^b = 25^b$, which is impossible.)

Analogously we obtain:

Proposition 6b. If $n \in \{3, 4, 5, \dots\}$ then $\left\{ \left(\sqrt{2 + 2/n} \right)^k \cdot (2/n)^l : k, l \in \{0, 1, 2, \dots\} \right\}$ is a dense subset of $(0, \infty)$.

Proposition 7. For each $n \in \{3, 4, 5, \dots\}$ there exists $k(n) \in \{0, 1, 2, \dots\}$ such that $1/2 \leq (2/n) \cdot \left(\sqrt{2 + 2/n} \right)^{k(n)} < 1$.

Proof. It is easy to check that

$$k(n) := \min \left\{ m \in \{0, 1, 2, \dots\} : 1/2 \leq (2/n) \cdot \left(\sqrt{2 + 2/n} \right)^m \right\}$$

satisfies our condition.

As a basic tool for our considerations we shall use the following lemma:

Lemma 1. Let $n \in \{2, 3, 4, \dots\}$ and $d, e \in D_n$ such that $2ne^2 > (n-1)d^2$, and define $r := \sqrt{4e^2 - 2\frac{n-1}{n}d^2}$. If there exists $\varepsilon \in D_n$ with $\varepsilon \leq 2r$, then $r \in D_n$.

Proof. We may assume that $\varepsilon \neq r$. Let $x, y \in \mathbb{R}^n$, $|x - y| = r$. According to [2], p.19, there exist points $p_1, \dots, p_n, \tilde{y}, \tilde{p}_1, \dots, \tilde{p}_n \in \mathbb{R}^n$ such that:

$$\begin{aligned}
|x - p_i| &= |y - p_i| = e \quad (1 \leq i \leq n), \\
|p_i - p_j| &= d \quad (1 \leq i < j \leq n), \\
|x - \tilde{y}| &= r, \\
|y - \tilde{y}| &= \varepsilon, \\
|x - \tilde{p}_i| &= |\tilde{y} - \tilde{p}_i| = e \quad (1 \leq i \leq n), \\
|\tilde{p}_i - \tilde{p}_j| &= d \quad (1 \leq i < j \leq n).
\end{aligned}$$

Let

$$S_{xy} := S_{y\tilde{y}} \cup \bigcup_{i=1}^n S_{xp_i} \cup \bigcup_{i=1}^n S_{yp_i} \cup \bigcup_{1 \leq i < j \leq n} S_{p_i p_j} \cup \bigcup_{i=1}^n S_{x\tilde{p}_i} \cup \bigcup_{i=1}^n S_{\tilde{y}\tilde{p}_i} \cup \bigcup_{1 \leq i < j \leq n} S_{\tilde{p}_i \tilde{p}_j}$$

and $f : S_{xy} \rightarrow \mathbb{C}^n$ preserves unit distance. Since

$$S_{xy} \supseteq \bigcup_{i=1}^n S_{xp_i} \cup \bigcup_{i=1}^n S_{yp_i} \cup \bigcup_{1 \leq i < j \leq n} S_{p_i p_j}$$

we conclude that f preserves the distances between x and p_i ($1 \leq i \leq n$), y and p_i ($1 \leq i \leq n$), and all distances between p_i and p_j ($1 \leq i < j \leq n$). Hence for all $1 \leq i \leq n$ $\varphi_n(f(x), f(p_i)) = \varphi_n(f(y), f(p_i)) = e^2$ and for all $1 \leq i < j \leq n$ $\varphi_n(f(p_i), f(p_j)) = d^2$. Since $S_{xy} \supseteq S_{y\tilde{y}}$ we conclude that $\varphi_n(f(y), f(\tilde{y})) = \varepsilon^2$. By Proposition 3 the Cayley-Menger determinant $\Delta(f(x), f(p_1), \dots, f(p_n), f(y))$ equals 0. Therefore

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & \dots & 1 & 1 & 1 \\ 1 & 0 & e^2 & e^2 & \dots & e^2 & e^2 & t \\ 1 & e^2 & 0 & d^2 & \dots & d^2 & d^2 & e^2 \\ 1 & e^2 & d^2 & 0 & \dots & d^2 & d^2 & e^2 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 1 & e^2 & d^2 & d^2 & \dots & 0 & d^2 & e^2 \\ 1 & e^2 & d^2 & d^2 & \dots & d^2 & 0 & e^2 \\ 1 & t & e^2 & e^2 & \dots & e^2 & e^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_n(f(x), f(y))$. Computing this determinant we obtain

$$(-1)^{n-1} \cdot d^{2n-2} \cdot t \cdot (nt + (2n-2)d^2 - 4ne^2) = 0.$$

Therefore

$$t = \varphi_n(f(x), f(y)) = \varphi_n(f(y), f(x)) = r^2$$

or

$$t = \varphi_n(f(x), f(y)) = \varphi_n(f(y), f(x)) = 0.$$

Analogously we may prove that

$$\varphi_n(f(x), f(\tilde{y})) = \varphi_n(f(\tilde{y}), f(x)) = r^2$$

or

$$\varphi_n(f(x), f(\tilde{y})) = \varphi_n(f(\tilde{y}), f(x)) = 0.$$

If $t = 0$ then the points $f(x)$ and $f(y)$ satisfy:

$$\begin{aligned} \varphi_n(f(x), f(x)) &= 0 = \varphi_n(f(y), f(x)), \\ \varphi_n(f(x), f(p_1)) &= e^2 = \varphi_n(f(y), f(p_1)), \\ &\dots \\ \varphi_n(f(x), f(p_n)) &= e^2 = \varphi_n(f(y), f(p_n)). \end{aligned}$$

By Proposition 4a the points $f(x), f(p_1), \dots, f(p_n)$ are affinely independent. Therefore by Proposition 5 $f(x) = f(y)$ and consequently

$$\varepsilon^2 = \varphi_n(f(y), f(\tilde{y})) = \varphi_n(f(x), f(\tilde{y})) \in \{r^2, 0\}.$$

Since $\varepsilon^2 \neq r^2$ and $\varepsilon^2 \neq 0$ we conclude that the case $t = 0$ cannot occur. This completes the proof of Lemma 1.

As corollaries we obtain:

Lemma 2. For $n \in \{2, 3, 4, \dots\}$, $d \in D_n$ and $m \in \{0, 1, 2, \dots\}$, $\left(\sqrt{2 + 2/n}\right)^m \cdot d \in D_n$.

Proof. For $e = \varepsilon = d$, Lemma 1 yields $r = \sqrt{2 + 2/n} \cdot d \in D_n$. This implies the assertion.

Lemma 3. Let $a, b \in D_2$ such that $a \leq \frac{4}{\sqrt{5}}b$. Then, $\sqrt{4b^2 - a^2} \in D_2$.

Proof. Since $a < 2b$, we have $r := \sqrt{4b^2 - a^2} > 0$. Now, for $\varepsilon := a$, from Lemma 1 we get the conclusion.

Lemma 4. For $d \in D_2$ and $l \in \{0, 1, 2, \dots\}$, $(\sqrt{3})^l \cdot d \in D_2$. Moreover, $\sqrt{11} \cdot d \in D_2$ and $5 \cdot d \in D_2$.

Proof. The first assertion follows from Lemma 2 and the second from Lemma 3 for $(a, b) := (d, \sqrt{3} \cdot d)$ and $(a, b) := (\sqrt{11} \cdot d, 3 \cdot d)$, respectively.

Additionally, for the plane case we need:

Lemma 5. If $d \in D_2$ then $(\sqrt{11}/5) \cdot d \in D_2$.

Proof. Let $d \in D_2$, $x, y \in \mathbb{R}^2$, $|x - y| = (\sqrt{11}/5) \cdot d$. Using the notation of Figure 1 we show that

$$S_{xy} := S_{y\tilde{y}} \cup \bigcup_{i=1}^2 S_{xp_i} \cup \bigcup_{i=1}^2 S_{yp_i} \cup S_{p_1p_2} \cup \bigcup_{i=1}^2 S_{x\tilde{p}_i} \cup \bigcup_{i=1}^2 S_{y\tilde{p}_i} \cup S_{\tilde{p}_1\tilde{p}_2}$$

where the sets corresponding to distances $\sqrt{3} \cdot d$, $\sqrt{11} \cdot d$ and $5 \cdot d$ are known to exist by Lemma 4.




Figure 1

$$|x - y| = |x - \tilde{y}| = (\sqrt{11}/5) \cdot d$$

$$|y - \tilde{y}| = d$$

$$|x - p_1| = |y - p_1| = |x - \tilde{p}_1| = |\tilde{y} - \tilde{p}_1| = \sqrt{11} \cdot d$$

$$|x - p_2| = |y - p_2| = |x - \tilde{p}_2| = |\tilde{y} - \tilde{p}_2| = \sqrt{3} \cdot d$$

$$|p_1 - p_2| = |\tilde{p}_1 - \tilde{p}_2| = 5 \cdot d$$

Assume that $f : S_{xy} \rightarrow \mathbb{C}^2$ preserves unit distance. Since

$$S_{xy} \supseteq S_{y\tilde{y}} \cup \bigcup_{i=1}^2 S_{xp_i} \cup \bigcup_{i=1}^2 S_{yp_i} \cup S_{p_1p_2}$$

we conclude that f preserves the distances between y and $\tilde{y}$, x and p_i ($i = 1, 2$), y and p_i ($i = 1, 2$), p_1 and p_2 . By Proposition 3, $\Delta(f(x), f(p_1), f(p_2), f(y))$ equals 0. Thus

$$\det \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 11d^2 & 3d^2 & t \\ 1 & 11d^2 & 0 & 25d^2 & 11d^2 \\ 1 & 3d^2 & 25d^2 & 0 & 3d^2 \\ 1 & t & 11d^2 & 3d^2 & 0 \end{bmatrix} = 0$$

where $t = \varphi_2(f(x), f(y))$. Computing this determinant we obtain

$$2d^2 \cdot t \cdot (11d^2 - 25t) = 0.$$

Therefore

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = \left(\left(\sqrt{11}/5 \right) \cdot d \right)^2$$

or

$$t = \varphi_2(f(x), f(y)) = \varphi_2(f(y), f(x)) = 0.$$

Analogously we may prove that

$$\varphi_2(f(x), f(\tilde{y})) = \varphi_2(f(\tilde{y}), f(x)) = \left(\left(\sqrt{11}/5 \right) \cdot d \right)^2$$

or

$$\varphi_2(f(x), f(\tilde{y})) = \varphi_2(f(\tilde{y}), f(x)) = 0.$$

If $t = 0$ then the points $f(x)$ and $f(y)$ satisfy:

$$\begin{aligned} \varphi_2(f(x), f(x)) &= 0 = \varphi_2(f(y), f(x)), \\ \varphi_2(f(x), f(p_1)) &= 11d^2 = \varphi_2(f(y), f(p_1)), \\ \varphi_2(f(x), f(p_2)) &= 3d^2 = \varphi_2(f(y), f(p_2)). \end{aligned}$$

By Proposition 4b the points $f(x), f(p_1), f(p_2)$ are affinely independent. Therefore by Proposition 5 $f(x) = f(y)$ and consequently

$$d^2 = \varphi_2(f(y), f(\tilde{y})) = \varphi_2(f(x), f(\tilde{y})) \in \left\{ \left(\left(\sqrt{11}/5 \right) \cdot d \right)^2, 0 \right\}.$$

Since $d^2 \neq \left(\left(\sqrt{11}/5 \right) \cdot d \right)^2$ and $d^2 \neq 0$ we conclude that the case $t = 0$ cannot occur. This completes the proof of Lemma 5.

Now we obtain:

Theorem 2a. If $x, y \in \mathbb{R}^2$ and $|x - y| = \left(\sqrt{11}/5 \right)^k \cdot \left(\sqrt{3} \right)^l$ (k, l are non-negative integers), then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^2$ such that each unit-distance preserving mapping from S_{xy} to $\mathbb{C}^2$ preserves the distance between x and y .

Proof. Since $1 \in D_2$, Lemmas 4 and 5 imply that

$$\left\{ \left(\sqrt{11}/5 \right)^k \cdot \left(\sqrt{3} \right)^l : k, l \in \{0, 1, 2, \dots\} \right\} \subseteq D_2.$$

Theorem 2b. If $x, y \in \mathbb{R}^n$ ($n \geq 3$) and $|x - y| = \left(\sqrt{2 + 2/n} \right)^k \cdot (2/n)^l$ (k, l are non-negative integers), then there exists a finite set S_{xy} with $\{x, y\} \subseteq S_{xy} \subseteq \mathbb{R}^n$ such that each unit-distance preserving mapping from S_{xy} to $\mathbb{C}^n$ preserves the distance between x and y .

Proof. By Lemma 2 if $d \in D_n$ ($n \geq 2$) then all distances $\left(\sqrt{2 + 2/n} \right)^m \cdot d$ ($m = 0, 1, 2, \dots$) belong to D_n . By Proposition 7 for each $n \in \{3, 4, 5, \dots\}$ there exists $k(n) \in \{0, 1, 2, \dots\}$ such that $1/2 \leq \rho(n) := (2/n) \cdot \left(\sqrt{2 + 2/n} \right)^{k(n)} < 1$. For $\varepsilon \in D_n$ we have $e := \left(\sqrt{2 + 2/n} \right)^{k(n)} \cdot \varepsilon \in D_n$ and $d := \sqrt{2 + 2/n} \cdot e \in D_n$. Therefore, by Lemma 1, if $\varepsilon \in D_n$, then $\rho(n) \cdot \varepsilon \in D_n$. Since $1 \in D_n$ we conclude that all distances $\rho(n)^m$ ($m = 0, 1, 2, \dots$) belong to D_n . For each $e \in D_n$ there exists $m \in \{0, 1, 2, \dots\}$ such that $\varepsilon := \rho(n)^m \leq (4/n) \cdot e$. Applying Lemma 1 for such ε and for $d := \sqrt{2 + 2/n} \cdot e$ we get:

(*) If $n \in \{3, 4, 5, \dots\}$ and $e \in D_n$, then $(2/n) \cdot e \in D_n$.

From Lemma 2 and (*) we obtain that for each non-negative integers k, l we have $\left(\sqrt{2 + 2/n} \right)^k \cdot (2/n)^l \in D_n$. This completes the proof of Theorem 2b.

Remark. If $n \in \{2, 3, 4, \dots\}$ and we define the set D_n for $f : S_{xy} \rightarrow \mathbb{R}^n$, then D_n is equal to the set of all positive algebraic numbers, see [11] and [12].

Finally we state:

Theorem 3. Each mapping $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ ($n \geq 2$) preserving unit distance preserves all distances i.e. satisfies

$$\forall x, y \in \mathbb{R}^n \quad \varphi_n(f(x), f(y)) = \varphi_n(x, y) = |x - y|^2,$$

provided that f is continuous with respect to the product topologies on $\mathbb{R}^n$ and $\mathbb{C}^n$.

Proof. Assume that $x, y \in \mathbb{R}^n$, $x \neq y$. Since $D_n \subseteq A_n$:

- (1) each map $f : \mathbb{R}^n \rightarrow \mathbb{C}^n$ preserving unit distance preserves all distances belonging to D_n i.e. for all $x, y \in \mathbb{R}^n$ if $|x - y| \in D_n$ then $\varphi_n(f(x), f(y)) = \varphi_n(x, y) = |x - y|^2$.

By Theorem 2a

$$\left\{ \left(\sqrt{11}/5 \right)^k \cdot \left(\sqrt{3} \right)^l : k, l \in \{0, 1, 2, \dots\} \right\} \subseteq D_2,$$

so by Proposition 6a D_2 is a dense subset of $(0, \infty)$. By Theorem 2b for all $n \geq 3$

$$\left\{ \left(\sqrt{2 + 2/n} \right)^k \cdot (2/n)^l : k, l \in \{0, 1, 2, \dots\} \right\} \subseteq D_n,$$

so by Proposition 6b for all $n \geq 3$ D_n is a dense subset of $(0, \infty)$. Therefore, for all $n \geq 2$:

- (2) there exists a sequence $\{y_k\} \subseteq \mathbb{R}^n$ tending to y such that for all k $|x - y_k| \in D_n$.

Since f and φ_n are continuous, (1) and (2) imply that

$$\varphi_n(f(x), f(y)) = \lim_{k \rightarrow \infty} \varphi_n(f(x), f(y_k)) = \lim_{k \rightarrow \infty} \varphi_n(x, y_k) = \varphi_n(x, y) = |x - y|^2.$$

Acknowledgement. The author wishes to thank the anonymous referee for the simplified proof of Theorem 2b and especially for her or his version of Theorem 2a with the proof. The author's original version of Theorem 2a states an analogical result for the distances $(2\sqrt{2}/3)^k \cdot (\sqrt{3})^l$ with non-negative integers k, l but the proof has been more complicated. The author also wishes to thank Professor Mowaffaq Hajja for improvement of the proof of Proposition 1.

References

- [1] F. S. BECKMAN AND D. A. QUARLES JR., *On isometries of euclidean spaces*, Proc. Amer. Math. Soc. 4 (1953), 810–815.
- [2] W. BENZ, *Geometrische Transformationen (unter besonderer Berücksichtigung der Lorentztransformationen)*, BI Wissenschaftsverlag, Mannheim, 1992.
- [3] L. M. BLUMENTHAL, *Theory and applications of distance geometry*, 2nd ed., Chelsea Publishing Co., New York, 1970.
- [4] K. BORSUK, *Multidimensional analytic geometry*, Polish Scientific Publishers, Warsaw, 1969.
- [5] N. BOURBAKI, *Éléments de mathématique, Livre II, Algèbre, Chapitre 4: Polynômes et fractions rationnelles, Chapitre 5: Corps commutatifs*, Deuxième édition, Hermann, Paris, 1959.
- [6] U. EVERLING, *Solution of the isometry problem stated by K. Ciesielski*, Math. Intelligencer 10 (1988), No. 4, p. 47.
- [7] D. J. H. GARLING, *A course in Galois theory*, Cambridge Univ. Press, Cambridge, 1995.
- [8] G. H. HARDY AND E. M. WRIGHT, *An introduction to the theory of numbers*, 5th ed., The Clarendon Press, Oxford University Press, New York, 1979.
- [9] M. KUCZMA, *An introduction to the theory of functional equations and inequalities: Cauchy's equation and Jensen's inequality*, Polish Scientific Publishers and Silesian University, Warsaw-Cracow-Katowice, 1985.
- [10] J. A. LESTER, *Distance preserving transformations*, in: 'Handbook of incidence geometry' (ed. F. Buekenhout), 921–944, North-Holland, Amsterdam, 1995.
- [11] A. TYSZKA, *Discrete versions of the Beckman-Quarles theorem*, Aequationes Math. 59 (2000), 124–133.
- [12] A. TYSZKA, *Discrete versions of the Beckman-Quarles theorem from the definability results of R. M. Robinson*, Algebra, Geometry & their Applications, Seminar Proceedings 1 (2001), 88–90, Yerevan State University Press.

- [13] P. B. YALE, *Automorphisms of the complex numbers*, Math. Mag. 39 (1966), 135–141.

A. Tyszka

Hugo Kołłątaj University

Technical Faculty

Balicka 104

PL-30-149 Kraków

Poland

e-mail: rttyszka@cyf-kr.edu.pl